

Errata for the Third Edition of Handbook of Differential Equations

LAST UPDATED: July 30, 2018

NOTES:

1. The latest errata is available from <http://www.mathtable.com/zwillinger/errata/>.
2. The home page for this book is <http://www.mathtable.com/hode/>.
3. You can reach the author at ZwillingerBooks@gmail.com.

I thank everyone who has contacted me about mistakes in this book!

1. Section 1, **Definition of Terms**, page 3. The commutator example may be misunderstood. The correction is to change

See Goldstein [6] for details.

To

Note that the “1”, as an operator, represents the identity. Hence, the first term is (in a different notation) $(xd)(1+d) = xd + xd^2$; it is not xd^2 . See Goldstein [6] for details.

(Thanks to David Goldsmith for this correction.)

2. Section 6, **Classification of Partial Differential Equations**, page 35.

- (a) The equation between equations (6.3) and (6.4) currently has the line:

$$u_{xy} = u_{\eta\eta}\eta_x\eta_y + \boxed{2}u_{\eta\zeta}(\eta_x\zeta_y + \eta_y\zeta_x) + u_{\zeta\zeta}\zeta_x\zeta_y + u_{\eta}\eta_{xy} + u_{\zeta}\zeta_{xy},$$

which is incorrect, it should have been:

$$u_{xy} = u_{\eta\eta}\eta_x\eta_y + u_{\eta\zeta}(\eta_x\zeta_y + \eta_y\zeta_x) + u_{\zeta\zeta}\zeta_x\zeta_y + u_{\eta}\eta_{xy} + u_{\zeta}\zeta_{xy},$$

- (b) The equation after equation (6.4) currently has the line:

$$\bar{B} = A\zeta_x\eta_x + B(\zeta_x\eta_y + \zeta_y\eta_x) + 2C\zeta_y\eta_y,$$

which is incorrect, it should have been (a “2” was missing)

$$\bar{B} = 2A\zeta_x\eta_x + B(\zeta_x\eta_y + \zeta_y\eta_x) + 2C\zeta_y\eta_y,$$

(Thanks to Hans Weertman for these corrections.)

3. Section 7, **Compatible Systems**, page 41, Special Case 3. The text for this special case is incorrect. It should be replaced with:

In the special case of $r = 1$, we have a system of m equations in m dependent variables. These equations do not require any side conditions.

(Thanks to Rusty Humphrey for this correction.)

4. Section 11, **Fixed Point Existence Theorems**, page 54

(a) The name “Schrauder” should be “Schauder”

(b) The following reference should be added:

J. SCHAUDER, “Der Fixpunktsatz in Funktionalräumen,” **Studia Math.**, 2, (1930), 171–180.

(Thanks to G. Friesecke for these corrections.)

5. Section 13, **Integrability of Systems**, page 65, Note number 11 contains “the sine–Gordan equation” when it should have “the sine–Gordon equation”.

(Thanks to Alain Moussiaux for this correction.)

6. Section 17, **Natural Boundary Conditions for a PDE**, page 77, The equation at the top of page 77, before equation (17.1) is now

$$J[\phi + h] - J[\phi] = \iint_R \left\{ L_{\phi_t} h_t + L_{\phi_{x_j}} h_{x_j} + L_{\phi} \right\} dt d\mathbf{x} + O(\|h\|^2),$$

This is incorrect, it should have been

$$J[\phi + h] - J[\phi] = \iint_R \left\{ L_{\phi_t} h_t + L_{\phi_{x_j}} h_{x_j} + L_{\phi} \boxed{h} \right\} dt d\mathbf{x} + O(\|h\|^2),$$

(Thanks to Zhuo Li for this correction.)

7. Section 27, **Canonical Forms**, page 118, reference number 2 is now

Bateman, H. *Partial Differential Equations of Mathematical Physics*, Dover Publications, New York, 1944.

Which is incorrect. The reference should have been

Bateman, H. *Differential Equations*, Longmans, Green and Co., New York, 1926, pages 75–79.

(Thanks to Ali Nejadmalayeri for this correction.)

8. Section 36, **Transformations of Second Order Linear ODEs – 1**, page 137, equation (36.6) is now $I(x) = (b - \frac{1}{4}a^2 - \frac{1}{2}\frac{da}{dx})$, which is incorrect. It should be $I(x) = (b + \frac{1}{4}a^2 - \frac{1}{2}\frac{da}{dx})$, (Thanks to Richard D. Rabbitt for this correction.)

9. Section 44.1.2, **Look-Up Technique**, page 169, the two equations

(a) Painlevé–Ince – modified

(b) Pinney

are both missing the “= 0” that should at the end of each.

(Thanks to Alain Moussiaux for these corrections.)

10. Section 44.1.3, **Look-Up Technique**, page 172, last equation before section 44.2, presently has

$$y^{(m)} = a \boxed{xy^{-m/2}}$$

This is incorrect, it should have been

$$y^{(m)} = a \boxed{yx^{-m/2}}$$

(Thanks to Flavio Noca for this correction.)

11. Section 50, **Clairaut’s Equation**, page 216, the equation between (50.5) and (50.6) is now

$$y''[2(xy' - \boxed{2})x - 2y'] = 0$$

which is incorrect. This expression should be

$$y''[2(xy' - y)x - 2y'] = 0$$

(Thanks to Bruno Muratori for this correction.)

12. Section 53, **Contact Transformation**, page 227, the second equation in equation 53.7 has the form

$$\dots = (2X^3 - 3X)^{1/3}$$

which is incorrect. This expression should be

$$\dots = \boxed{C} (2X^3 - 3X)^{1/3}$$

(Thanks to Alain Moussiaux for this correction.)

13. Section 72, **Green's functions**, page 292, From above equation (72.9) to that equation the text is presently:

Using the second method, we find the eigenvalues and eigenfunctions to be

$$\lambda_n = \frac{n\pi}{L}, \quad \phi_n(x) = \sin \lambda_n x = \sin\left(\frac{n\pi x}{L}\right),$$

so that

$$G(x; z) = \frac{2L}{n\pi} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{n\pi z}{L}\right).$$

which is incorrect; the text should have been

Using the second method, we find the eigenvalues and eigenfunctions to be

$$\lambda_n = \left(\frac{n\pi}{L}\right)^2, \quad \phi_n(x) = \sin \lambda_n x = \sin\left(\frac{n\pi x}{L}\right),$$

so that

$$G(x; z) = \sum_{n=1}^{\infty} \left[\frac{2L^2}{n^2\pi^2} \right] \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{n\pi z}{L}\right).$$

(Thanks to Luis Alberto Fernandez for this observation.)

14. Section 79, **Integrating Functions**, page 326, note number 10, the following should be added:

The general solution to $u_x = yu_y$ is $u = f(x + \log y)$, where f is an arbitrary function.

(Thanks to Alain Moussiaux for this observation.)

15. Section 80, **Interchanging Dependent and Independent Variables**, page 327,

- (a) In Example 3, the nonlinear equation is given as " $y''(x - y)y'^3$ ", which is incorrect. It should have been " $y''(y - x)y'^3$ ".

(Thanks to Alain Moussiaux for this correction.)

- (b) In Note number 2, the reference to Bender and Orszag should be section 1.5, not 1.6.

(Thanks to James Dare for this correction.)

- (c) A better citation for reference number 3 is: McAllister, B. L. and Thorne, C.J. "Reverse differential equations and others that can be solved exactly", *Studies Appl. Math*, 6, 1952.

(Thanks to Daniele Ritelli for this correction.)

16. Section 85, **Reduction of order**, page 354, note number 2 presently contains

More generally, if $\{z_1(x), \dots, z_p(x)\}$ are linearly independent solutions of equation (85.6), then the substitution

$$y(x) = \begin{bmatrix} z_1 & \dots & z_p & v \\ z'_1 & \dots & z'_p & v' \\ \vdots & & \vdots & \vdots \\ z_1^{(p)} & \dots & z_p^{(p)} & v^{(p)} \end{bmatrix}$$

reduces equation (85.7) to a linear ordinary differential equation of order $n - p$ for $v(x)$.

This should be changed to

More generally, if $\{z_1(x), \dots, z_p(x)\}$ are linearly independent solutions of equation (85.6), then the substitution

$$y(x) = \begin{bmatrix} z_1 & \dots & z_p & z \\ z'_1 & \dots & z'_p & z' \\ \vdots & & \vdots & \vdots \\ z_1^{(p)} & \dots & z_p^{(p)} & z^{(p)} \end{bmatrix} \phi(x) \quad (1)$$

where $\phi(x)$ need not be specified, reduces equation (85.6) to a linear ordinary differential equation of order $n - p$ for $y(x)$. The following explains why.

With the above, $y(x)$ can be written in the form

$$y(x) = A(x)z^{(p)} + B(x)z^{(p-1)} + \dots, \quad A(x) \neq 0$$

and its derivatives have the form

$$y'(x) = A(x)z^{(p+1)} + \dots, \quad y''(x) = A(x)z^{(p+2)} + \dots,$$

These equations can be used to eliminate $\{z^{(p)}, \dots, z^{(n)}\}$ and (85.6) will take the form

$$b_0 y^{(n-p)} + \dots + b_{n-p} y + V = 0 \quad (2)$$

where V is linear in the $\{z, z', \dots, z^{(p-1)}\}$.

We argue that $V \equiv 0$ as follows: Consider equation (2) as a differential equation of degree $p - 1$ in z (via the V term). If $z = z_i$ (for any $i = 1, 2, \dots, p$) then $y = 0$ from equation (1). Hence, from equation (2) it must be that $V|_{z=z_i} = 0$. Hence $\{z_i\}_{i=1,2,\dots,p}$ is a collection of p linearly independent solutions to a differential equation of degree $p - 1$; possible only if $V \equiv 0$.

(Thanks to Unal Goktas for this correction.)

17. Section 87, **Matrix Riccati Equations**, page 358. The second line in equation (87.4) is now

$$\frac{dy}{dt} = b(t)(y^2 - x^2) - 2a(t)xy \boxed{-} 2cy$$

Which is incorrect, it should have been

$$\frac{dy}{dt} = b(t)(y^2 - x^2) - 2a(t)xy + 2cy$$

(Thanks to both Peter Sherwood and Alain Moussiaux for this correction.)

18. Section 93, **Superposition**, page 373, the last line contains the equation

$$L[y] = y'' + a(x)y' + b(x) = f(x)$$

Which is incorrect. This should have been

$$L[y] = y'' + a(x)y' + b(x) \boxed{y} = f(x)$$

(Thanks to Young Kim for this correction.)

19. Section 96, **Vector Ordinary Differential Equations** pages 384-385, In note number 9 the second equation is incorrect. All the text after “Alternately, if the ...” should be deleted.

(Thanks to Frankie Liu for this correction.)

20. Section 106, **Inverse Scattering**, page 416, the **Applicable to** statement should have at the end

having the form of (106.2)

(Thanks to G. Friesecke for this observation.)

21. Section 106, **Inverse Scattering**, page 418, Note number 5 gives a Lax pair for the equation $u_t + u_{xx} - 2uu_x = 0$, which is not quite the Burger’s equation. (Notice the minus sign before the last term.)

(Thanks to Bruno Muratori for this correction.)

22. Section 118, **Chaplygin's Method**, page 465, equations (118.5) and (118.6) and the surrounding text are now

Then define $u_1(x)$ to be the solution of

$$y' = M(x)y + N(x), \quad y(x_0) = y_0. \quad (118.5)$$

and define $v_1(x)$ to be the solution of

$$y' = \widehat{M}(x)y + \widehat{N}(x), \quad y(x_0) = y_0. \quad (118.6)$$

Which is incorrect. This should have been (note that the definitions have been switched):

Then define $v_1(x)$ to be the solution of

$$y' = M(x)y + N(x), \quad y(x_0) = y_0. \quad (118.5)$$

and define $u_1(x)$ to be the solution of

$$y' = \widehat{M}(x)y + \widehat{N}(x), \quad y(x_0) = y_0. \quad (118.6)$$

(Thanks to Bruno Van der Bossche for these corrections.)

23. Section 123, **Graphical Analysis: The Phase Plane**, pages 479, 480.

(a) In the text for example 1 it says

... The curve figure 123.2 is given by $\text{determinant} = (\text{trace})^2$; only centers can occur along this curve.

which is incorrect; it should have said

... The curve in figure 123.2 is given by $\text{determinant} = (\text{trace}/2)^2$. Centers occur along the curve defined by $\text{trace} = 0$.

(b)

(Thanks to Zhuo Li for these corrections.)

24. Section 136, **Monge's Method**, pages 523–524,

(a) Equation (136.5) contains, in part

$$\dots = \boxed{\frac{\partial z}{\partial y}} + 6y$$

which is incorrect. This expression should be

$$\dots = \boxed{\frac{\partial z}{\partial x}} + 6y$$

(b) Equation (136.10) contains, in part

$$\dots + \psi(2\boxed{z} + y^2)$$

which is incorrect. This expression should be

$$\dots + \psi(2\boxed{x} + y^2)$$

(Thanks to Alain Moussiaux for this correction.)

25. Section 139, **Perturbation Method: Method of Averaging**, pages 532–533,

(a) In equations (139.3) and (139.5) the last “cos” in each case should be a “sin”.

(b) The two equations in (139.9) are each missing a final closing parenthesis.

(Thanks to Gerald Teschl for these corrections.)

26. Section 143, **Perturbation Method: Regular Perturbation**, page 554, equations (143.5 b) and (143.7 b) both have “ $y_1(0) = 1$ ” which is incorrect; they should have been “ $y_1(0) = 0$ ”.

(Thanks to Frank Scharf for this corrections.)

27. Section 145, **Picard Iteration**, page 561, note number one, the following should be added:

However, the successive approximations are guaranteed to converge to the true solution for all x sufficiently close to zero provided f is a continuously differentiable function.

(Thanks to G. Friesecke for this observation.)

28. Section 148, **Soliton-Type Solutions**, pages 567–569,

(a) In equation (148.3) the term cv_ζ should be $-cv_\zeta$.

(b) In equation (148.4) the term $(v_\zeta)^2$ should be $\frac{1}{2}(v_\zeta)^2$.

(c) An additional note should be added on page 569 to state

With the standard choice of $A = B = 0$, the solution to (148.4) can be solved in terms of elementary functions:

$$v(x) = \frac{3c}{\sigma} \left(\operatorname{sech} \left(\frac{\sqrt{cx}}{2} \right) \right)^2$$

(Thanks to G. Friesecke for these corrections.)

29. Section 180, [Runge–Kutta Methods](#), pages 691, 696

(a) Equation (180.3) is missing some “ h ” terms. Presently there is:

$$\begin{aligned} k_1 &= f(x_0, y_0), \\ k_2 &= f(x_0 + \frac{1}{2}h, y_0 + \frac{1}{2}k_1), \\ k_3 &= f(x_0 + \frac{1}{2}h, y_0 + \frac{1}{2}k_2), \\ k_4 &= f(x_0 + h, y_0 + k_3). \end{aligned} \tag{3}$$

which is incorrect. It should have been:

$$\begin{aligned} k_1 &= f(x_0, y_0), \\ k_2 &= f(x_0 + \frac{1}{2}h, y_0 + \frac{1}{2}\boxed{h}k_1), \\ k_3 &= f(x_0 + \frac{1}{2}h, y_0 + \frac{1}{2}\boxed{h}k_2), \\ k_4 &= f(x_0 + h, y_0 + \boxed{h}k_3). \end{aligned} \tag{4}$$

(b) Note number 9 is incorrect and should be deleted.

30. Section 172, [Pseudospectral Method](#), page 772, presently has:

$$\left. \frac{\partial u}{\partial x} \right|_{x=x_k} \simeq \frac{1}{3h}(u_{k+1} - u_{k-1}) - \frac{1}{6h}(u_{k+2} - u_{k-2}).$$

and

$$\left. \frac{\partial u}{\partial x} \right|_{x=x_k} \simeq \frac{1}{2h}(u_{k+1} - u_{k-1}) - \frac{1}{3h}(u_{k+2} - u_{k-2}) + \frac{1}{30h}(u_{k+3} - u_{k-3}).$$

and

$$\left. \frac{\partial u}{\partial x} \right|_{x=x_k} = \sum_{j=1}^{\infty} \frac{2(-1)^{j+1}}{jh}(u_{k+j} - u_{k-j}).$$

Which are all incorrect. They should have been:

$$\left. \frac{\partial u}{\partial x} \right|_{x=x_k} \simeq \frac{2}{3h}(u_{k+1} - u_{k-1}) - \frac{1}{12h}(u_{k+2} - u_{k-2}).$$

and

$$\left. \frac{\partial u}{\partial x} \right|_{x=x_k} \simeq \frac{3}{4h}(u_{k+1} - u_{k-1}) - \frac{3}{20h}(u_{k+2} - u_{k-2}) + \frac{1}{60h}(u_{k+3} - u_{k-3}).$$

and

$$\left. \frac{\partial u}{\partial x} \right|_{x=x_k} = \sum_{j=1}^{\infty} \frac{(-1)^{j+1}}{jh}(u_{k+j} - u_{k-j}).$$

(Thanks to Didier Clamond for these corrections.)

NEW MATERIAL

If Dan Zwillinger were writing *Handbook of Differential Equations* today, additional sections that would be added include:

- Definition and Concepts

1. Classification of Equations

- (a) Roman Kozlov, *The group classification of a scalar stochastic differential equation*, Journal of Physics A: Mathematical and Theoretical, Volume 43, Number 5, 12 January 2010, <http://iopscience.iop.org/article/10.1088/1751-8113/43/5/055202/meta>

Lie's group classification of the stochastic differential equation $dx = f(t, x)dt + g(t, x)dW(t)$

Group dimension	Basis operators	Equation
0	No symmetries	$dx = f(t, x)dt + dW(t)$
1	$X_1 = \frac{\partial}{\partial t}$	$dx = f(x)dt + dW(t)$
2	$X_1 = \frac{\partial}{\partial t}, \quad X_2 = 2t\frac{\partial}{\partial t} + x\frac{\partial}{\partial x}$	$dx = \frac{\alpha}{x}dt + dW(t), \quad \alpha \neq 0$
3	$X_1 = \frac{\partial}{\partial t}, \quad X_2 = 2t\frac{\partial}{\partial t} + x\frac{\partial}{\partial x}, \quad X_3 = \frac{\partial}{\partial x}$	$dx = dW(t)$

- Exact Methods

1. Differential Constraints

- (a) Oleg V Kaptsov and Alexey V Schmidt, *Linear determining equations, differential constraints and invariant solutions*, 31 Aug 2003, <https://arxiv.org/pdf/math-ph/0309001.pdf>
- (b) Boris Kruglikov, *Symmetry approaches for reductions of PDEs, differential constraints and Lagrange-Charpit method*, <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.520.5511&rep=rep1&type=pdf>
- (c) Peter J. Olver, *Direct Reduction and Differential Constraints*, http://www-users.math.umn.edu/~olver/s_/dc.pdf

2. Eigenvalue problems

- (a) Donald S. Cohen, *An Integral Transform Associated with Boundary Conditions Containing an Eigenvalue Parameter*, SIAM J. Appl. Math., Volume 14, Number 5, pages 1164–1175, <https://epubs.siam.org/doi/abs/10.1137/0114093>.

3. Fokas Method

- (a) See Wikipedia entry https://en.wikipedia.org/wiki/Fokas_method
- (b) Bernard Deconinck, Thomas Trogdon, and Vishal Vasan, *The Method of Fokas for Solving Linear Partial Differential Equations*, SIAM Review, Volume 56, Number 1, pages 159–186, <https://epubs.siam.org/doi/abs/10.1137/110821871>

4. Specialized Techniques

- (a) Barton L. Willis, *An Extensible Differential Equation Solver*, ACM SIGSAM Bulletin, Volume 35, Number 1, March 2001, <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.577.5712&rep=rep1&type=pdf>
 - (b) J F Carinena, J Grabowski, and J de Lucas, *Superposition rules for higher order systems and their applications*, Journal of Physics A: Mathematical and Theoretical Volume 45, Number 18, 20 April 2012, <http://iopscience.iop.org/article/10.1088/1751-8113/45/18/185202/meta>
 - (c) Gromov’s approach
 - Deane Yang, *Gromov’s Approach to Solving Underdetermined Systems of PDE’s*, http://www.deaneyang.com/papers/underdetermined_ode.pdf
 - Victor Lehenkyi, *The Integrability of Some Underdetermined Systems*, Proceedings of Institute of Mathematics of NAS of Ukraine, 2000, Volume 30, Part 1, 157–164, <https://pdfs.semanticscholar.org/bab0/ebb721ae26a91d517e2ad5277583e6569.pdf>
- Approximate Methods
 - 1. **Renormalization Group Method**
 - (a) Hayato Chiba, *Simplified Renormalization Group Method for Ordinary Differential Equations*, <http://www2.math.kyushu-u.ac.jp/~chiba/paper/rg2.pdf>
 - (b) R. E. Lee DeVille, Anthony Harkin, Matt Holzer, Kresimir Josic, and Tasso J. Kaper, *Analysis of a Renormalization Group Method for Solving Perturbed Ordinary Differential Equations*, 14 August 2006, <https://faculty.math.illinois.edu/~rdeville/research/FinalRGpaper.pdf>
 - (c) Eleftherios Kirkinis, *The Renormalization Group: A Perturbation Method for the Graduate Curriculum*, SIAM REVIEW, Volume 54, Number 2, pages 374–388 <https://epubs.siam.org/doi/pdf/10.1137/080731967>
 - Numerical Methods
 - 1. **Block Operators**
 - (a) Jared L. Aurentz and Lloyd N. Trefethen, “Block Operators and Spectral Discretizations”, *SIAM Review*, Vol. 39, No. 2, pages 423–446, <https://epubs.siam.org/doi/abs/10.1137/16M1065975>
 - 2. **Parallel Computer Algorithms**
 - (a) Terry Haut and Beth Wingate, *An Asymptotic Parallel-in-Time Method for Highly Oscillatory PDEs*, SIAM J. Sci. Comput., Volume 36, Number 2, A693–A713, <https://epubs.siam.org/doi/abs/10.1137/130914577>
 - (b) Marcos Rodriguez, Fernando Blesa, and Barrio, Roberto, *OpenCL parallel integration of ordinary differential equations: Applications in computational dynamics*, Computer Physics Communications, 192, (July 2015).
 - 3. **Poincare Maps**
 - (a) See Wikipedia entry https://en.wikipedia.org/wiki/Poincare_map

- (b) Warwick Tucker, *Computing accurate Poincaré maps*, Physica D 171 (2002) 127–137, <http://www2.math.uu.se/~warwick/main/papers/accuratePoincare.pdf>

4. Quantum Algorithms

- (a) Dominic W. Berry, Andrew M. Childs, Aaron Ostrander, and Guoming Wang, *Quantum algorithm for linear differential equations with exponentially improved dependence on precision*, 17 Feb 2017, <https://arxiv.org/pdf/1701.03684.pdf>
- (b) Dominic W. Berry, *High-order quantum algorithm for solving linear differential equations*, 28 Jan 2014, <https://arxiv.org/pdf/1010.2745.pdf>.

UPDATED MATERIAL

There have been advances in many fields that should be included in an updated version of *Handbook of Differential Equations*. There are also references that were previously overlooked.

- **5. Chaos in Dynamical Systems**, add the references

- Monica A. Garcia–Nustes and Jorge A. Gonzalez, *Universal functions and exactly solvable chaotic systems*, 7 Nov 2008, <https://arxiv.org/pdf/0811.1179.pdf>.
- J. C. Sprott, *A simple chaotic delay differential equation*, Physics Letters A 366 (2007), pages 397–402, <http://sprott.physics.wisc.edu/pubs/paper304.pdf>
- Y. Charles Li, *Chaos in Partial Differential Equations*, 4 September 2009, <https://arxiv.org/pdf/0909.0910.pdf>.

- **6. Classification of Partial Differential Equations**

Add the reference

- M. S. Chong, A. E. Perry, and B. J. Cantwell, *A general classification of three-dimensional flow fields*, Physics of Fluids, A 2(5), 1990, pages 765–777.

Add the new **Example 4**

- Consider the 3-by-3 system $\frac{dx}{dt} = Ax$, where A is constant. The eigenvalues of A satisfy $\lambda^3 + P\lambda^2 + Q\lambda + R = 0$ where

$$P = -\text{trace}(A)$$

$$Q = \frac{1}{2}(P^2 - \text{trace } A^2)$$

$$R = -\det A$$

The surface S , defined by $27R^2 + (4P^3 - 18PQ)R + (4Q^3 - P^2Q^2) = 0$, divides the real solutions from the complex solutions. When $\{P, Q, R\}$ are real, the surface S can be split up into two surfaces S_{1a} and S_{1b} given by $R = R_a$ and $R = R_b$ where:

$$R_a = \frac{1}{3}P \left(Q - \frac{2}{9}P^2 \right) - \frac{2}{27}(P^2 - 3Q)^{3/2}$$

$$R_b = \frac{1}{3}P \left(Q - \frac{2}{9}P^2 \right) + \frac{2}{27}(P^2 - 3Q)^{3/2}$$

The classification of solution trajectories in (P, Q, R) space is:

1. node / node / node
 - (a) nodes are stable if: $[P > 0, 0 < Q < \frac{P^2}{3}, 0 < R < R_b]$
 - (b) nodes are unstable if: $[P < 0, 0 < Q < \frac{P^2}{3}, R_a < R < 0]$
2. node / node / star node
 - (a) nodes are stable if: $[P > 0, \frac{P^2}{4} < Q < \frac{P^2}{3}, R = R_a]$ or $[P > 0, 0 < Q < \frac{P^2}{3}, R = R_b]$
 - (b) nodes are unstable if: $[P < 0, \frac{P^2}{4} < Q < \frac{P^2}{3}, R = R_a]$ or $[P < 0, 0 < Q < \frac{P^2}{3}, R = R_b]$
3. star node / star node / star node
 - (a) nodes are stable if: $[P > 0, Q = \frac{P^2}{3}, R = R_a = R_b = \frac{P^3}{27}]$
 - (b) nodes are unstable if: $[P < 0, Q = \frac{P^2}{3}, R = R_a = R_b = \frac{P^3}{27}]$
4. line node-saddle / line node-saddle / node
 - (a) nodes are stable if: $[P > 0, 0 < Q < \frac{P^2}{4}, R = 0]$
 - (b) nodes are unstable if: $[P < 0, 0 < Q < \frac{P^2}{4}, R = 0]$
5. line node-saddle / line node-saddle / star node
 - (a) nodes are stable if: $[P > 0, Q = \frac{P^2}{4}, R = R_a = 0]$
 - (b) nodes are unstable if: $[P < 0, Q = \frac{P^2}{4}, R = R_b = 0]$
6. node / saddle / saddle
 - (a) node is stable if: $[P \geq 0, Q < \frac{P^2}{4}, R_a < R < 0]$ or $[P < 0, Q < 0, R_a < R < 0]$
 - (b) node is unstable if: $[P \leq 0, Q < \frac{P^2}{4}, 0 < R < R_a]$ or $[P > 0, Q < 0, 0 < R < R_b]$
7. star node / saddle / saddle
 - (a) node is stable if: $[P \geq 0, Q < \frac{P^2}{4}, R = R_a]$ or $[P < 0, Q < 0, R = R_a]$
 - (b) node is unstable if: $[P \leq 0, Q < \frac{P^2}{4}, R = R_b]$ or $[P > 0, Q < 0, R = R_b]$
8. line node-saddle / line node-saddle / no flow
 - (a) both line node-saddles are stable if: $[P > 0, Q = 0, R = 0]$
 - (b) both line node-saddles are unstable if: $[P < 0, Q = 0, R = 0]$
 - (c) one line node-saddle is stable and the other unstable if: [for all $P, Q < 0, R = 0]$
9. focus / stretching
 - (a) focus is stable if: $[P \geq 0, Q > \frac{P^2}{4}, R < 0]$ or $[P \geq 0, Q < \frac{P^2}{4}, R < R_a]$ or $[P < 0, Q > 0, R < PQ]$ or $[P < 0, Q > 0, R < R_a]$
 - (b) focus is unstable if: $[P < 0, Q > \frac{P^2}{3}, PQ < R < 0]$ or $[P < 0, \frac{P^2}{4} < Q < \frac{P^2}{3}, R_b < R < 0]$ or $[P < 0, 0 < Q < \frac{P^2}{3}, PQ < R < R_a]$
10. focus / compressing
 - (a) focus is stable if: $[P > 0, Q > \frac{P^2}{4}, 0 < R < PQ]$ or $[P > 0, \frac{P^2}{4} < Q < \frac{P^2}{3}, 0 < R < R_a]$ or $[P > 0, 0 < Q < \frac{P^2}{3}, R_b < R < PQ]$
 - (b) focus is unstable if: $[P \leq 0, Q > \frac{P^2}{4}, R > 0]$ or $[P \leq 0, Q < \frac{P^2}{4}, R < R_b]$ or $[P > 0, Q > 0, R < PQ]$ or $[P > 0, Q > 0, R < R_a]$
11. focus / no flow
 - (a) focus is stable if: $[P > 0, Q > \frac{P^2}{4}, R = 0]$
 - (b) focus is unstable if: $[P < 0, Q > \frac{P^2}{4}, R = 0]$
12. center
 - (a) stretching if $[P < 0, Q > 0, R = PQ]$
 - (b) compressing if $[P > 0, Q > 0, R = PQ]$

(c) no flow if $[P = 0, Q > 0, R = 0]$

• **8. Conservation Laws**, add the references

- J. J. H. Bashingwa and A. H. Kara, *A basis of hierarchy of generalized symmetries and their conservation laws for the (3+1)-dimensional diffusion equation*, 6 Dec 2017, <https://arxiv.org/pdf/1712.02117.pdf>.
- Wen-Xiu Ma, *Conservation laws by symmetries and adjoint symmetries*, 11 June 2017, <https://arxiv.org/pdf/1707.03496.pdf>.
- Roman O. Popovych and Alexander Bihlo, *Inverse problem on conservation laws*, 9 May 2017, <https://arxiv.org/pdf/1705.03547.pdf>.

• **84. Pfaffian Differential Equations**, add the reference

- David Delphenich, *The role of integrability in a large class of physical system*, 16 March 2017, <https://arxiv.org/ftp/arxiv/papers/1210/1210.4976.pdf>.

• **66. Factoring Operators**

Add the reference

- Mark Giesbrecht, Albert Heinle, and Viktor Levandovskyy, *Factoring Differential Operators in n Variables*, 31 March 2014, <https://arxiv.org/pdf/1404.0002.pdf>.

Add the following

The paper by Giesbrecht *et al.*, presents an algorithm for factoring partial differential operators for a large class of operators. Technically they find factorizations over many algebras, including the polynomial n^{th} Weyl algebra. The n^{th} Weyl algebra Q_n is defined by

$$Q_n := \mathbb{K} \langle x_1, \dots, x_n, \partial_1, \dots, \partial_n \rangle$$

$$\partial_i x_j = \begin{cases} x_j \partial_i & \text{if } i \neq j \\ q_i x_j \partial_i + 1 & \text{if } i = j \end{cases}$$

$$0 = \partial_i \partial_j - \partial_j \partial_i = x_i x_j - x_j x_i$$

for $i, j = 1, \dots, n$ where the $\{q_1, \dots, q_n\}$ are units in \mathbb{K} . A computer algebra implementation of their algorithm is available in SINGULAR.

Examples in the paper include:

1. $\partial^2 = \partial \partial = \left(\partial + \frac{1}{x+c} \right) \left(\partial - \frac{1}{x+c} \right)$
2. $\partial^3 - x \partial - 2 = \left(\partial + \frac{1}{x} \right) \left(\partial^2 - \frac{1}{x} \partial - x \right)$
3. $x (\partial^3 - x \partial - 2) = \partial (x \partial^2 - x^2 - \partial)$
4. $(\partial_1 + 1)^2 (\partial_1 + x_1 \partial_2) = (x_1 \partial_1 \partial_2 + \partial_1^2 + x_1 \partial_2 + 2 \partial_2) (\partial_2 + 1)$.

Additionally, using their package,

- 60 distinct factorizations are found for $x_1 x_2^2 x_3^2 \partial_1 \partial_2^2 + x_2 x_3^3 \partial_2$
- 60 distinct factorizations are found for $(x_1^2 \partial_1 + x_1 x_2 \partial_2) (\partial_1 \partial_2 + \partial_1^2 \partial_2^2 x_1 x_2)$

– 12 distinct factorizations are found for $(x_1^4 - 1)x_1\partial_1^2 + (1 + 7x_1^4)\partial_1 + 8x_1^3$.

• **95. Variation of parameters**

Add the following as note number 5:

Explicit solutions using variation of parameters:

1. If the linear second order equation $L[y] = y'' + P_1(x)y' + P_0(x)y = R(x)$ has the homogeneous solutions $y_1(x)$ and $y_2(x)$ (i.e., $L[y_i] = 0$), then the solution to the original equation may be written as

$$\begin{aligned} y(x) &= -y_1(x) \int \frac{y_2(x)R(x)}{W(y_1, y_2)} dx + y_2(x) \int \frac{y_1(x)R(x)}{W(y_1, y_2)} dx, \\ &= y_1(x) \int \frac{\begin{vmatrix} 0 & y_2 \\ R & y_2' \end{vmatrix}}{W(y_1, y_2)} dx + y_2(x) \int \frac{\begin{vmatrix} y_1 & 0 \\ y_1' & R \end{vmatrix}}{W(y_1, y_2)} dx \end{aligned}$$

where $W(y_1, y_2) = y_1y_2' - y_1'y_2 = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$ is the Wronskian.

2. If the linear third order equation $L[y] = y''' + P_2(x)y'' + P_1(x)y' + P_0(x)y = R(x)$ has the homogeneous solutions $y_1(x)$, $y_2(x)$, and $y_3(x)$ (i.e., $L[y_i] = 0$), then the solution to the original equation may be written as

$$\begin{aligned} y(x) &= y_1(x) \int \frac{\begin{vmatrix} 0 & y_2 & y_3 \\ 0 & y_2' & y_3' \\ R & y_2'' & y_3'' \end{vmatrix}}{W(y_1, y_2, y_3)} dx + y_2(x) \int \frac{\begin{vmatrix} y_1 & 0 & y_3 \\ y_1' & 0 & y_3' \\ y_1'' & R & y_3'' \end{vmatrix}}{W(y_1, y_2, y_3)} dx \\ &\quad + y_3(x) \int \frac{\begin{vmatrix} y_1 & y_2 & 0 \\ y_1' & y_2' & 0 \\ y_1'' & y_2'' & R \end{vmatrix}}{W(y_1, y_2, y_3)} dx \end{aligned} \tag{5}$$

where $W(y_1, y_2, y_3) = \begin{vmatrix} y_1 & y_2 & y_3 \\ y_1' & y_2' & y_3' \\ y_1'' & y_2'' & y_3'' \end{vmatrix}$ is the Wronskian.

• **112. Separation of Variables**, add the references

- H. S. Cohl and H. Volkmer, *Separation of variables in an asymmetric cyclidic coordinate system*, 16 Jan 2013, <https://arxiv.org/pdf/1301.3559.pdf>.
- H. S. Cohl and H. Volkmer, *Expansions for a fundamental solution of Laplace's equation on R^3 in 5-cyclidic harmonics*, 14 Nov 2013, <https://arxiv.org/pdf/1311.3514.pdf>.
- George Pogosyan, Alexey Sissakian, and Pavel Winternitz, *Separation of Variables and Lie Algebra Contractions. Applications to Special Functions*, 8 Oct 2003, <https://arxiv.org/pdf/math-ph/0310011.pdf>.
- A. Szereszewski and A. Sym, *On Darboux's approach to R-separability of variables. Classification of conformally flat 4-dimensional binary metrics*, Journal of Physics A: Mathematical and Theoretical, Volume 48, Number 38, 25 August 2015 <http://iopscience.iop.org/article/10.1088/1751-8113/48/38/385201/meta>.

- **128 Interval Analysis**, add the reference
 - Fahimeh Goodarzi, Mahmoud Hadizadeh, and Farideh Ghoreishi, *An interval solution for the n -th order linear ODEs with interval initial conditions*, Mathematical Communications, Volume 18, Number 1, 2013, <http://www.mathos.unios.hr/mc/index.php/mc/article/view/242>.
- **156. Available Software**, add the references
 - FreeFem++ – a numerical PDE solver – <http://www.freefem.org/>
 - JuliaDiffEq – a numerical differential equations solver for Julia – <http://juliadiffeq.org/packages.html>
 - deSolve – a numerical ODE/ADE/DDE solver in R – <http://cran.ms.unimelb.edu.au/web/packages/deSolve/vignettes/deSolve.pdf> – <https://cran.r-project.org/web/packages/deSolve/index.html>
 - SUNDIALS: SUite of Nonlinear and Differential/ALgebraic Equation Solvers – <https://computation.llnl.gov/projects/sundials>
- **182. Integrating Stochastic Equations**, add the references
 - Desmond J. Higham, *An Algorithmic Introduction to Numerical Simulation of Stochastic Differential Equations*, SIAM REVIEW, Volume 43, Number 3, 2001, pages 525–546, <https://epubs.siam.org/doi/pdf/10.1137/S0036144500378302>
 - Cian O. Mahony, *The Numerical Analysis of Stochastic Differential Equations*, 8 September 2006, <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.117.8043&rep=rep1&type=pdf>
 - Kevin Burrage and Pamela Burrage, *High Strong Order Explicit Runge-Kutta Methods for Stochastic Ordinary Differential Equations*, Applied Numerical Mathematics, 22 (1–3), October 1996, <https://www.sciencedirect.com/science/article/pii/S016892749600027X>